

BOSTON PUBLIC LIBRARY



3 9999 06317 706 5

152.

COMPARATIVE DIETARY TOXICITIES OF PESTICIDES TO BIRDS

MAY 21 1972

DEPT. OF THE INTERIOR

UNITED STATES DEPARTMENT OF THE INTERIOR
FISH AND WILDLIFE SERVICE
BUREAU OF SPORT FISHERIES AND WILDLIFE

Special Scientific Report—Wildlife No. 152

UNITED STATES DEPARTMENT OF THE INTERIOR, ROGERS C. B. MORTON, *SECRETARY*
Nathaniel P. Reed, *Assistant Secretary for Fish and Wildlife and Parks*
Fish and Wildlife Service
Bureau of Sport Fisheries and Wildlife, Spencer H. Smith, *Director* (Acting)

COMPARATIVE DIETARY TOXICITIES OF PESTICIDES TO BIRDS

By

Robert G. Heath
James W. Spann
Elwood F. Hill
James F. Kreitzer

Patuxent Wildlife Research Center
Division of Wildlife Research
Bureau of Sport Fisheries and Wildlife
Laurel, Maryland 20810



Special Scientific Report--Wildlife No. 152
Washington, D. C. • February 1972

For sale by the Superintendent of Documents, U.S. Government Printing Office
Washington, D.C. 20402 - Price 60 cents
, Stock Number 2410-0314

Table of Contents

	<u>Page</u>
ABSTRACT	iv
INTRODUCTION	1
PROCEDURES	1
Protocol for Determining Lethal Dietary Concentrations of Chemicals	2
The Probit Analysis	3
The Dieldrin Standard	3
Rationale for the Protocol and the LC_{50}	3
TOXICITY STATISTICS	5
Relative Toxicities	5
Estimating LC's for the General Percentage of Response . .	6
Utility of Toxicity Statistics	6
DISCUSSION	7
Toxicity Comparisons	7
Comparisons of Species Sensitivity	9
The 8-day Mortality Pattern	9
ACKNOWLEDGMENTS	10
REFERENCES	10
Table 1	13
Table 2	22
Table 3	46
Table 4	54
APPENDIX 1	56
APPENDIX 2	57

ABSTRACT

This report presents measurements of the lethal dietary toxicity of 89 pesticidal chemicals to young bobwhites, Japanese quail, ring-necked pheasants, and mallards. Toxicity is expressed as the median lethal concentration (LC_{50}) of active chemical in a 5-day ad libitum diet. LC_{50} 's and associated statistics are derived by methods of probit analysis. Endrin consistently was the most toxic chemical while aldrin and dieldrin were among the six most toxic chemicals of those tested on all species. In general, organophosphates were less toxic than aldrin or dieldrin, and herbicides were of a low order of toxicity. There were obvious inconsistencies in the relative sensitivity of the four species to various chemicals.

INTRODUCTION

It is now well documented in scientific literature that pesticidal contamination of ecosystems can alter the status of animal populations through diverse, often complex modes of action (see Stickel, 1968). Direct lethal toxicity is the most obvious mode of action, and laboratory measurements of lethality are basic in predicting the immediate impact of pesticides in the environment.

The following report presents measurements of the lethal toxicities of 89 pesticidal chemicals as administered for 5 days in diets of young birds of four species: bobwhite (*Colinus virginianus*), Japanese quail (*Coturnix coturnix japonica*), ring-necked pheasant (*Phasianus colchicus*), and mallard (*Anas platyrhynchos*). All work was conducted at the Patuxent Wildlife Research Center. The study was designed to provide statistically reliable estimates of the relative toxicity of any two chemicals as tested. It also permits comparisons of species sensitivity to pesticides in the diet.

None of the findings is intended to imply chemical safety beyond the scope of the study; while measurements of lethal toxicity constitute an important first step in evaluating pesticidal hazard, comprehensive estimates of total population effects require knowledge of a variety of parameters (physical, chemical, biochemical, and toxicological), involving suspect degradation products as well as parent chemicals. Such elusive, delayed effects as reproductive impairment or alterations of critical behavior patterns may be highly detrimental. Further, there is little apparent correspondence between a chemical's lethal toxicity and its capacity to induce sublethal complications. For example, DDE, a ubiquitous metabolite of DDT, is not especially toxic but produces serious reproductive effects in various species of birds (Heath et al., 1969; Wiemeyer and Porter, 1970).

PROCEDURES

All test birds were incubator-hatched progeny of breeding colonies maintained on the Patuxent Center. Bobwhites, pheasants, and mallards were phenotypically indistinguishable from wild birds; our Japanese quail colony was started from eggs obtained in 1964 from Auburn University, Auburn, Ala. All colonies were randomly outbred so that findings, although probably associated with greater variances than those among inbred strains, can be more readily related to wild populations. Tests were conducted with bobwhites, pheasants, and mallards during spring and summer; Japanese quail were tested throughout the remainder of the year.

Protocol for Determining Lethal Dietary Concentrations of Chemicals:

The protocol is essentially that described by Heath and Stickel (1965) for testing dietary toxicity of pesticides to birds, and proposed by the U.S. Department of the Interior for use in pesticide registration (ref. 19). Toxicity is expressed as a median lethal concentration (LC_{50}) of chemical in dry diet. The LC_{50} is defined herein as ppm toxicant in an ad libitum diet expected to produce 50 percent mortality among 2- to 3-week-old birds in 8 days comprising 5 days on treated diet followed by 3 days on untreated diet. The final 3 days are included to detect chemical mortality induced beyond the dosage period; otherwise LC_{50} 's would tend to be overestimated. All measurements are in terms of the active ingredient, exclusive of diluents or impurities.

Tests with gallinaceous chicks were conducted in laboratory-housed brooder units. Each unit consists of six separated tiers of four wire pens in which heat is thermostatically controlled. Tests with mallard ducklings were conducted in weatherproof wooden pens on straw-covered concrete slabs, each supplied with an infra-red heating lamp and a trough of running water.

Birds were randomly assigned to study pens on the day preceeding a test and acclimated on untreated commercial diet prior to dosage. Usually 10 birds were assigned per pen, although six to 15 birds were sometimes used, depending upon availability. Dosage was never initiated before birds were 9 days old, to avoid possible interference of chemical intake by yolk sac absorption and to exclude hatchling mortality. There was no attempt to sex the birds at this early age.

To prepare test diets, chemical was dissolved in corn oil or, as necessary, in propylene glycol, and was then mixed thoroughly with dry commercial mash in a ratio of 2 parts of solution to 98 parts of feed by weight. An equal amount of pure corn oil was added to "control" diets. Thorough mixing of feed and solution was accomplished with a commercial food mixer.

Each chemical was generally administered in six dietary concentrations spaced geometrically over a span intended to produce mortality ranging from 10 to 90 percent during the 8-day period. Concentrations were selected after range finding with three widely spaced levels. One pen of birds was used per concentration, and facilities were adequate to test as many as seven chemicals (i.e., 42 concentrations) simultaneously. Included with each set of chemicals was a dieldrin "standard" of six concentrations, and three to six pens of control birds fed untreated diets. A completely randomized design was used in each experiment, with treatments and birds randomly assigned to pens.

The Probit Analysis:

Deaths in each pen were recorded daily, and total deaths during the 8-day period were used to derive percentages of mortality resulting from each dietary concentration of chemical. There were infrequent deaths among the untreated controls, and these were used to adjust the data for extraneous mortality by means of Abbott's formula (Finney, 1952).

All LC_{50} 's and associated statistics were derived by methods of probit analysis as described by Finney (1952) and programmed for computer by Daum and Killcreas (1966). The program calculates a number of maximum likelihood statistics including: the 95 percent confidence limits of each LC_{50} ; the slope, and its standard deviation, of the weighted linear regression of probits on log-concentration; and the relative toxicity, with confidence limits, of any two chemicals after testing regression lines for parallelism and heterogeneity. The program fits up to 10 parallel probit regression lines simultaneously; thus chemicals tested in any one experiment were analysed as a set.

The Dieldrin Standard:

A dieldrin standard was used in every experiment, i.e., with every set of chemicals tested simultaneously. The necessity for a standard to adjust comparisons of toxicity of two or more chemicals tested at different times or locations has been discussed by various authors (Sun, 1950; Finney, 1952; Bliss, 1952). Without a standard, differences in animal sensitivity between experiments, whether due to physiological or environmental causes, can lead to biased comparisons of toxicity. Comparisons adjusted through a standard will be unbiased providing any differences in sensitivity between studies affect the toxicity estimates of the standard in the same proportion as those of the test chemicals. We selected dieldrin as a standard because it is a well-known chemical that consistently provided an acceptable probit regression line. DDT was used initially but tended to give heterogeneous results.

Not all chemicals are sufficiently toxic to be lethal at concentrations that might reasonably be expected in the environment. We therefore established a ceiling of 5,000 ppm for an LC_{50} , with occasional exceptions. Chemicals shown in range finding tests to have LC_{50} 's greater than 5,000 ppm were not tested further.

Rationale for the Protocol and the LC_{50} :

When the protocol was designed, several investigators had reported that the lethality of a pesticide mixed in the diet could differ markedly from that of the concentrate administered as a single

oral dose via capsule or gavage (Stickel et al., 1965). In human toxicology, where there is major concern with accidental or suicidal poisoning from sudden doses of concentrate, the classic single oral dose is clearly appropriate. Our particular interest, however, was in dietary toxicity because ingestion is undoubtedly the predominant route of exposure in wild species.

The primary objective in designing the protocol was to establish a dosage period short enough to approach acute exposure, yet long enough to insure adequate feeding activity by all test birds to produce sufficient mortality for meaningful determinations. The 5-day period was selected and, we believe, has proved to be satisfactory. (The important subject of chronicity, involving rates of chemical storage and excretion as well as degradation, is best studied using dosage periods longer than 5 days [Hayes, 1967; Weil and McCollister, 1963].)

The administration of toxicant blended in the diet provides in a sense, an "applied" measurement of toxicity, in that incorporated in each measurement are not only the digestive factors that determine uptake of chemical from the feed into the system, but also the propensity of the animal to ingest food containing the pesticide. Therefore, we have expressed lethality in terms of "ppm chemical in feed" rather than attempting to convert to a ratio of unit of chemical per unit of body weight, such as mg/kg.

Further reasons made such conversions impractical. Food consumption could not be measured accurately, since feed scattered by birds became mixed with litter and droppings, and its weight could only be estimated. Each bird in a given pen actually ingested a slightly different amount of toxicant, and there is no assurance that if each bird had consumed the average mg/kg for the pen the same percentage of mortality would have resulted. A conversion to mg/kg/day would provide a better measurement than mg/kg but would not overcome the problem of food spillage. At best it would be a rough measurement based on pen averages and would take additional time to compute. For these reasons conversions were not attempted.

Proper measurements of mg/kg/day require that birds be caged individually, weighed daily, their dosage predetermined and diet prepared daily, and a procedure utilized to insure complete ingestion of the preparation. Stress caused by excessive handling might conceivably lead to anomalous results.

Food consumption, corrected for estimated spillage, was measured to detect reductions in intake due to treatment. Repellency was rarely demonstrated (as substantiated by satisfactory probit regression lines) except occasionally at the higher dosage levels. There was some reduction in average daily consumption associated with high mortality, probably due more to intoxication than repellency.

TOXICITY STATISTICS

An alphabetical listing of common plus chemical names of compounds tested comprises table 1. Detailed toxicity statistics for each chemical and species are given in table 2, while table 3 ranks chemicals by order of toxicity for each species. Table 4 compares species sensitivity to several classes of compounds. Detailed descriptions are included with each table.

Relative Toxicities:

The term "relative toxicity," as used in this study, is the ratio of the dietary concentrations of two chemicals expected to produce a given percentage of mortality, as tested, in a specified population of birds. For example, if twice the concentration of chemical B is needed to produce the same percentage of mortality as a given concentration of chemical A, then chemical A has twice the lethal toxicity of chemical B. In general, the relative toxicity of chemical A to chemical B (at a given level of mortality) is the required concentration of B divided by that of A.

The relative toxicity of two chemicals may be expressed unconditionally as the ratio of their LC_{50} 's provided (1) the level of tolerance of the test subjects did not differ between the experiments in which the chemicals were tested, and (2) the two probit regression lines are parallel. The first condition can be assumed valid only for chemicals tested in the same completely randomized experiment. However, adjustments for tolerance differences between experiments can be made through the ratio of the LC_{50} 's of the respective dieldrin standards. The second condition, parallelism, is assumed if the slopes of the regression lines are not shown to be different at a specified level of statistical significance. If the lines are not parallel, relative toxicity will vary with dietary concentration and therefore cannot be expressed as a constant.

The relative toxicity of dieldrin (RTD) to each chemical tested is presented for each species in table 2, with a notation if parallelism was rejected. To calculate the relative toxicities of other pairs of chemicals, adjusted through the dieldrin standard, we propose using the data in table 2 according to the procedure presented in appendix 1.

There is perhaps a tendency to suppose that if one chemical is, say, twice as toxic as another, it will cause twice the mortality at a given dosage. Such a relationship does not hold true, however, as can readily be demonstrated by constructing two hypothetical dosage-mortality lines on log-probability paper. The valid comparison of toxicity is between the amounts of chemicals required to produce the same effect, and not between the percentages of mortality resulting from equal dosages.

Estimating LC's for the General Percentage of Response:

Lethal concentrations for percentages of mortality other than the median can be estimated from the data in table 2. The computations involve transforming the LC_{50} to its common logarithm and the desired percentage of mortality to its probit, the probit of 50 percent being 5. If we let k equal the new percentage of mortality for which we wish to estimate the lethal dietary concentration (i.e., the LC_k), and b equal the particular slope value from table 2, then

$$\log LC_k = (5 - \text{probit } k) / b - \log LC_{50}$$

The antilog of $\log LC_k$ is, of course, the desired estimate. Tables for transforming percentages to probits can be found in various statistical texts, including Finney (1952).

In this study, there appeared to be little advantage to estimating response dosages other than the LC_{50} . It can be estimated more precisely than any other value, per given effort, and serves to provide efficient estimates of relative toxicity. At times, estimates of extreme values, such as the LC_5 or LC_{99} , may be important; however, such estimates are best determined from especially designed experiments, since extrapolation from a standard probit regression line can be misleading if the true regression equation has some curvature (Finney, 1952).

Utility of Toxicity Statistics:

The immediate utility of the toxicity statistics lies, perhaps, in their providing readily calculable comparisons of the short-term dietary toxicity of any two of the chemicals to a given species. Further, the slopes of the probit regression lines indicate the rate of increase in lethal hazard resulting from a proportional increase in exposure, i.e., the steeper the slope, the more rapid the increase in lethality.

Often, pesticidal contamination in the natural diet can be measured by chemical analysis of field samples. When this is possible, the degree of hazard may be predicted by direct comparison of ppm chemical in the natural diet with the lethality of dietary concentrations in laboratory studies.

When residues in the natural diet cannot be determined, short-term hazard may be estimated, as mentioned by Tucker and Crabtree (1970), by comparing the test chemical (T) with a chemical (K) known to produce mortality in the field when applied at a given rate. For example, if in the laboratory T is twice as toxic as K (i.e., the LC_{50} of T is one-half that of K), then we should expect T to produce mortality in the field if applied at one-half or more the

stipulated rate for K. Expressions of hazard that relate quantities of chemical lethal in laboratory doses to quantities of chemical applied per unit area (say, per square foot) have been suggested in the literature. The approach is necessarily a manipulation of the one immediately above; and while convenient, it cannot offer greater accuracy. Clearly, the accuracy of all such predictions will depend upon how representative the laboratory measurements are of toxicity in the field.

Multiple, interacting factors (physical, chemical, biochemical, and toxicological) determine the total hazard of a pesticide once released into the environment, as outlined in detail by Kenaga (1968). Thus findings herein should not be extrapolated beyond the limits of the protocol without caution and qualification.

DISCUSSION

Toxicity Comparisons:

More than 4500 intra-species estimates of relative toxicity can be derived from the data in table 2 in addition to those involving the dieldrin standard. Although it would be impractical to list them in this report, certain general comparisons should be noted.

Endrin was consistently the most toxic chemical as tested. It was virtually twice as toxic as any other chemical with the exception of DRC-1339 (Starlicide), which was tested only on Japanese quail. Dieldrin and the closely related aldrin were always among the six most toxic chemicals of those tested on all species, and to pheasants they were the second and third most toxic chemicals.

Organophosphates generally proved to be less toxic in the diet than aldrin or dieldrin. Of 22 organophosphates tested on bobwhite, only three (phosphamidon, fenthion, and Dasanit) were more toxic; of 23 tested on Japanese quail, only parathion and methyl parathion were more toxic; and of 23 tested on mallards, only Dasanit and diazinon were more toxic. None of 21 organophosphates was more toxic to pheasants than aldrin or dieldrin.

Data from Tucker and Crabtree (1970), who present a variety of estimates of acute toxicity using encapsulated doses, show that when chemical concentrate is administered as a single oral dose, organophosphate pesticides are generally more toxic to mallards and pheasants than either aldrin or dieldrin. Limited data for Japanese quail suggest the same tendency, but there are too few comparable data for bobwhite.

We can only theorize about the differences due to the two methodologies. Apparently organophosphate degradation in treated

feed prior to consumption was not a factor: virtually identical results were obtained when Dasanit and demeton concentrates were prepared and mixed in feed daily and when the diets were prepared at the onset of the 5-day period. It has been reported, however, that absorption of some organochlorine compounds through the gastrointestinal wall is more efficient when chemical is blended in the diet than when administered as a single dose of concentrate (Stickel et al., 1965). With organophosphate compounds we doubt that this difference is pronounced. Moreover, exposure tends to be gradual when toxicant is dispursed throughout the diet, often giving the system time to degrade the relatively unstable organophosphate compounds but not the notably stable organochlorine compounds.

Dietary LC_{50} 's for certain of the chemicals have been estimated previously. Those for six Aroclor products (polychlorinated biphenyl mixtures) given by Heath et al. (1970) were derived from the same data used herein, and slight changes represent computerized refinements. The same is true for several organochlorine pesticides, excepting technical DDT, presented by Heath et al. (1965); however, DDT estimates are derived from subsequent tests, and differences appear to be largely the result of a change in source of technical DDT. Further, poor reproducibility (heterogeneity) of DDT results may have contributed to the differences.

Gill et al. (1970) have given 5-day dietary LC_{50} 's of technical and p,p'-DDT, p,p'-DDD, and p,p'-DDE for pheasant chicks. We have tested technical DDT, a technical DDD marketed as Rhothane, and p,p'-DDE, always in the same completely randomized experiment for a given species. Gill's LC_{50} 's of DDD and DDE (522 and 1086 ppm) corresponded quite well with ours for pheasants (579 and 841 ppm), but theirs of technical DDT (935) is higher than we present here (311 ppm). We suspect a difference between the technical DDT formulations, since our technical DDT was consistently more toxic than either Rhothane or DDE to all species. Their LC_{50} of 550 ppm for p,p'-DDT corresponds better to ours for technical DDT.

DDE is generally reported to be much less toxic than DDD; and although it is probably the most abundant pesticide residue in the environment, DDE is thought to present little hazard in a lethal sense. With the exception of pheasants, we found this toxicity relationship not to hold true. Testing both chemicals simultaneously in completely randomized experiments, we derived the following LC_{50} 's for DDE and DDD: bobwhite, 825 and 2178 ppm; Japanese quail, 1355 and 3165 ppm; and mallards, 3572 and 4814 ppm. Thus DDE was about 2 1/2 times as toxic as DDD to both bobwhite and Japanese quail and almost 1 1/2 times as toxic to mallards.

There appears to be a definite reversal in the toxicity of diazinon and Guthion to birds compared to mammals. Both in dietary and in encapsulated doses (Tucker and Crabtree, 1970) diazinon was from seven to 38 times as toxic as Guthion to mallards and pheasants; however, oral LD_{50} 's for rats (Gaines, 1960 and 1969) show Guthion to be at least 19 times as toxic as diazinon.

Herbicides were generally of a low order of toxicity. Only diquat and paraquat consistently produced mortality, the latter being the most toxic herbicide tested. Otherwise most LC_{50} 's were greater than 5000 ppm, and in many tests there was no mortality at that concentration.

Comparisons of Species Sensitivity:

There were obvious inconsistencies in the relative sensitivity of the four species to various chemicals, as is shown in table 4. Mallards generally tolerated the highest dietary concentrations, but to toxaphene, terepene polychlorinates, and Ceresan M, the mallard was the most sensitive species. Bobwhites, Japanese quail, and pheasants were similar in overall sensitivity to organochlorine compounds; but to most organophosphates, bobwhites and Japanese quail were more sensitive than pheasants. There was a major difference in the sensitivity of bobwhites and Japanese quail to the Aroclors, bobwhites being the most sensitive of the four species and Japanese quail the least sensitive, as reported earlier (Heath et al., 1970); otherwise there was no discernible difference in the overall sensitivity of the two species.

Such inconsistencies in relative sensitivity among four species, especially when three are gallinacious species, strongly suggest that certain ones not tested will be much more sensitive to the different chemicals. Studies using mature house sparrows, blue jays, cardinals, and bobwhites (Hill, 1971) also show considerable variation in species sensitivity.

The 8-Day Mortality Pattern:

Death rarely occurred during the first day of dosage and only occasionally on the second day. Mortality tended to accelerate from the third through fifth day of dosage, subsided on the first day of untreated diet (i.e., the sixth day), and deaths rarely occurred thereafter. The pattern of mortality was generally similar for all effective dosages, the level of mortality being a function of a particular dosage.

Two very toxic chemicals - Ceresan M and Dasanit - produced notable exceptions to this pattern, especially in tests with mallards. Ceresan M produced little mortality during the 5 days of dosage, although the birds had become severely intoxicated by the fifth day. Mortality began on the first day of untreated diet and was so heavy during the second and third days that mortality on untreated diet was recorded for an additional six days. Only one of eight birds survived at 60 ppm (active ingredient), and none at levels of 74 ppm and higher. There were no deaths, however, at 20 ppm or among controls.

The response to Dasanit was completely opposite to that of Ceresan M. Virtually all mortality occurred during the first day, the extent of mortality depending upon dosage level; if a bird survived the initial day of dosage, it apparently became tolerant of the chemical, regardless of dosage level, and survived. As shown earlier, chemical breakdown in feed was definitely not a factor.

ACKNOWLEDGMENTS

A number of individuals made significant contributions to the study. Dr. L. F. Stickel was codeveloper of the protocol, and Dr. J. L. Buckley was instrumental in its development. Mr. W. H. Stickel and Dr. E. H. Dustman provided valuable information regarding the status of new pesticides; and the Center's chemists, especially Messrs. W. L. Reichel, E. Cromartie, and Dr. R. M. Prouty, assisted with various chemical problems. Mr. C. Vance was in charge of aviculture, and Messrs. N. H. Kruhm, C. Mills, and F. T. Polak serviced test facilities. Mrs. H. L. Young conducted the computer analyses and assisted in compiling tables, and Mrs. H. M. Nelson performed preliminary computations. We wish to thank the various chemical companies that provided technical grade compounds used in the study.

REFERENCES

1. Bliss, C. I.
1952. The Statistics of Bioassay. (Reprinted, with additions, from Vitamin Methods, Vol. II.) Academic Press, New York.
Pp. 445-628.
2. Daum, R. J., and W. Killcreas.
1966. Two computer programs for probit analysis. Bulletin of the Entomological Society of America 12 (4): 365-369.
3. Finney, D. J.
1952. Probit Analysis, 2d ed. Cambridge University Press.
318 p.
4. Gaines, T. B.
1960. The acute toxicity of pesticides to rats. Toxicology and Applied Pharmacology 2 (1): 88-99.
5. _____
1969. Acute toxicity of pesticides. Toxicology and Applied Pharmacology 14 (3): 515-534.
6. Gill, J. A., B. J. Verts, and A. G. Christensen.
1970. Toxicities of DDE and some other analogs of DDT to pheasants. Journal of Wildlife Management 34 (1): 223-226.

7. Hayes, W. J., Jr.
1967. The 90-dose LD₅₀ and a chronicity factor as measures of toxicity. *Toxicology and Applied Pharmacology* 11 (2): 327-335.
8. Heath, R. G., J. W. Spann, and J. F. Kreitzer.
1969. Marked DDE impairment of mallard reproduction in controlled studies. *Nature* 224 (5214): 47-48.
9. _____, _____, _____, and C. Vance.
1970. Effects of polychlorinated biphenyls on birds. *Proceedings XV International Ornithological Congress, The Hague, 30 August - 5 September, 1970. Edited K. H. Voous. E. J. Brill Publishers, Leiden.*
10. _____, _____, and C. Vance.
1965. Pen studies of avian toxicology at Patuxent. P. 23-24 in U.S. Fish and Wildlife Circular 226.
11. _____, and L. F. Stickel.
1965. Protocol for testing the acute and relative toxicity of pesticides to birds. P. 18-21 in U.S. Fish and Wildlife Circular 226.
12. Hill, E. F.
1971. Toxicity of selected mosquito larvicides to some common avian species. *Journal of Wildlife Management* 35 (4): 757-762.
13. Kenaga, E. E.
1968. Guidelines for evaluating the properties of pesticides for safe use in the wildlife environment. *Down to Earth* 23 (4): 11-18.
14. Snedecor, G. W., and W. G. Cochran.
1967. *Statistical Methods*, 6th ed. Iowa State Univ. Press, Ames. 593 p.
15. Stickel, L. F.
1968. Organochlorine pesticides in the environment. U.S. Bureau of Sport Fisheries and Wildlife, Special Scientific Report -- Wildlife No. 119. 32 p.
16. Stickel, W. H., W. E. Dodge, W. G. Sheldon, J. B. DeWitt, and L. F. Stickel.
1965. Body condition and response to pesticides in woodcocks. *Journal of Wildlife Management* 29 (1): 147-155.

17. Sun, Yun-Pei.
1950. Toxicity index -- an improved method of comparing the relative toxicity of insecticides. Journal of Economic Entomology 43 (1): 45-53.
18. Tucker, R. K., and D. G. Crabtree.
1970. Handbook of toxicity of pesticides to wildlife. U.S. Bureau of Sport Fisheries and Wildlife, Resource Publication No. 84. 131 p.
19. U.S. Fish and Wildlife Service.
1964. Procedures for evaluation of toxicity of pesticides to wildlife. Pesticides Review Staff. 77 p.
20. Weil, C. S., and D. D. McCollister.
1963. Relationship between short-and long-term feeding studies in designing an effective toxicity test. Journal of Agricultural and Food Chemistry 11 (6): 486-491.
21. Wiemeyer, S. N., and R. D. Porter.
1970. DDE thins eggshells of captive American kestrels. Nature 227 (5259): 737-738.

Table 1 provides an alphabetical listing of official coined (common) names and principal registered names of the chemicals tested. Chemical name, class, and primary usage are shown with the common name, or with the registered name where a common name does not exist. Other registered names refer the reader to the common name: Chemical names follow the nomenclature of *Chemical Abstracts*.

Table 1. Common and chemical names of pesticidal compounds tested in 5-day diets of second-week birds

Common or registered name	Chemical name	Class ^a	Use
Abate	0,0,0',0'-tetramethyl 0,0'-thiodi- <u>p</u> -phenylene phosphorothioate	OP	insecticide
Accothion	(see fenitrothion)		
aldrin	1,2,3,4,10,10-hexachloro-1,4,4a,5, 8,8a-hexahydro-1,4- <u>endo-exo</u> -5,8- dimethanonaphthalene	OC	insecticide
aminotriazole	(see amitrole)		
amitrole	3-amino-1,2,4-triazole	O	herbicide
Aramite	2-(<u>p</u> -tert-butylphenoxy) isopropyl- 2'-chloroethyl sulfite	O	acaricide
Aroclor 1221	polychlorinated biphenyl (21% chlorine)	OC	industrial
Aroclor 1232	polychlorinated biphenyl (32% chlorine)	OC	industrial
Aroclor 1242	polychlorinated biphenyl (42% chlorine)	OC	industrial
Aroclor 1248	polychlorinated biphenyl (48% chlorine)	OC	industrial
Aroclor 1254	polychlorinated biphenyl (54% chlorine)	OC	industrial
Aroclor 1260	polychlorinated biphenyl (60% chlorine)	OC	industrial
Aroclor 1262	polychlorinated biphenyl (62% chlorine)	OC	industrial
atrazine	2-chloro-4-ethylamino-6- isopropylamino- <u>s</u> -triazine	O	herbicide
Baygon	<u>o</u> -isopropoxyphenyl methyl carbamate	C	insecticide

Table 1. Common and chemical names of pesticidal compounds tested in 5-day diets of second-week birds--continued

Common or registered name	Chemical name	Class ^a	Use
Baytex	(see fenthion)		
Cadminate	cadmium succinate 60% (cadmium 29%)	OM	fungicide
captan	N-trichloromethylthio-4-cyclohexene-1,2-dicarboximide	O	fungicide
carbaryl	1-naphthyl N-methylcarbamate	C	insecticide
Casoron	(see dichlobenil)		
Ceresan M	N-(ethylmercuri)-p-toluene sulfonanilide	OM	fungicide
chlordan	1,2,4,5,6,7,8,8-octachloro-2,3,3a,4,7,7a-hexahydro-4,7-methanoindene	OC	insecticide
Co-Ral	0,0-diethyl 0-3-chloro-4-methyl-1-oxo-2H-1-benzopyran-7-yl phosphorothioate	OP	insecticide
Cygon	(see dimethoate)		
2,4-D, acetamide	2,4-dichlorophenoxyacetic acid, acetamide	PH	herbicide
2,4-D, butoxy-ethanol ester	2,4-dichlorophenoxyacetic acid, butoxyethanol ester	PH	herbicide
2,4-D, dimethyl-amine salt	2,4-dichlorophenoxyacetic acid, dimethylamine salt	PH	herbicide
dalapon	2,2-dichloropropionic acid, sodium salt	O	herbicide
2,4-DB	4-(2-dichlorophenoxy) butyric acid	PH	herbicide
Dasanit	0,0-diethyl 0-p-(methylsulfinyl) phenyl phosphorothioate	OP	insecticide

Table 1. Common and chemical names of pesticidal compounds tested in 5-day diets of second-week birds--continued

Common or registered name	Chemical name	Class ^a	Use
DDD	1,1-dichloro-2,2- <u>bis</u> -(p-chlorophenyl) ethane	OC	insecticide
DDE	1,1-dichloro-2,2- <u>bis</u> -(p-chlorophenyl) ethylene	OC	(degradation product of DDT)
DDT	1,1,1-trichloro-2,2- <u>bis</u> -(p-chlorophenyl) ethane	OC	insecticide
DDVP	2,2-dichlorovinyl dimethyl phosphate	OP	insecticide
Delnav	(see dioxathion)		
demeton	0,0-diethyl-0-2-(ethylthio) ethyl phosphorothioate and 0,0-diethyl S-2-(ethylthio) ethyl phosphorothioate	OP	insecticide acaricide
diazinon	0,0-diethyl 0-(2-isopropyl 4-methyl-6-pyrimidinyl) phosphorothioate	OP	insecticide
Dibrom	(see naled)		
dichlobenil	2-6-dichlorobenzonitrile	O	herbicide
dichlone	2,3-dichloro-1,4-naphthoquinone	O	fungicide
dichlorobenzophenone	4,4'-dichlorobenzophenone	OC	(degradation product of DDT)
dicofol	4,4'-dichloro- <u>alpha</u> -trichloromethylbenzhydrol	OC	acaricide
dieldrin	1,2,3,4,10,10-hexachloro-6,7-epoxy-1,4,4a,5,6,7,8,8a-octahydro-1,4- <u>endo-exo</u> -5,8-dimethanonaphthalene	OC	insecticide
dimethoate	0,0-dimethyl S-(N-methyl-carbamoylmethyl) phosphorodithioate	OP	insecticide acaricide
dioxathion	2,3- <u>p</u> -dioxane S,S-bis (0,0-diethylphosphorodithioate)	OP	acaricide insecticide

Table 1. Common and chemical names of pesticidal compounds tested in 5-day diets of second-week birds--continued

Common or registered name	Chemical name	Class ^a	Use
Dipterex	(see trichlorfon)		
diquat	1,1'-ethylene-2,2'-dipyridinium dibromide	O	herbicide desicant
disulfoton	O,O-diethyl S-2-(ethylthio) ethyl phosphorodithioate	OP	insecticide acaricide systemic
DiSyston	(see disulfoton)		
Dithane D-14	(see nabam)		
diuron	3-(3,4-dichlorophenyl)-1,1-dimethylurea	U	herbicide
DRC-1339	3-chloro- <u>p</u> -toluidine hydrochloride	OC	avicide
Dursban	O,O diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate	OP	insecticide
Dyfonate	O-ethyl-S-phenyl-ethylphosphono-dithioate	OP	insecticide
Dylox	(see trichlorfon)		
endosulfan	6,7,8,9,10,10-hexachloro-1,5,5a,6,9,9a-hexahydro-6,9-methano-2,4,3-benzodioxathiepin-3-oxide	OC	insecticide
endrin	1,2,3,4,10,10-hexachloro-6,7-epoxy-1,4,4a,5,6,7,8,8a-octa-hydro-1,4- <u>endo-endo</u> -5,8-dimethanonaphthalene	OC	insecticide
EPN	O-ethyl O- <u>p</u> -nitrophenyl phenylphosphonothioate	OP	insecticide acaricide
fenac	2,3,6-trichlorophenyl-acetic acid	O	herbicide
fenitrothion	O,O-dimethyl O-(4-nitro- <u>m</u> -tolyl) phosphorothioate	OP	insecticide
fenthion	O,O-dimethyl O-[4-(methylthio)- <u>m</u> -toyl] phosphorothioate	OP	insecticide

Table 1. Common and chemical names of pesticidal compounds tested in 5-day diets of second-week birds--continued

Common or registered name	Chemical name	Class ^a	Use
fenuron	3-phenyl-1,1-dimethylurea	U	herbicide
Guthion	<u>O,O</u> -dimethyl S-[4-oxo-1,2,3-benzotriazin-3(4H)-ylmethyl] phosphorodithioate	OP	insecticide
heptachlor	1,4,5,6,7,8,8-heptachloro-3a, 4,7,7a-tetrahydro-4,7- <u>endo</u> -methanoindene	OC	insecticide
Imidan	0,0-dimethyl S-phthalimidomethyl phosphorodithioate	OP	insecticide acaricide
Kelthane	(see dicofol)		
Lannate	(see methomyl)		
lindane	Gamma isomer of 1,2,3,4,5,6-hexachlorocyclohexane of 99+% purity	OC	insecticide
malathion	S-[1,2,-bis(ethoxycarbonyl)ethyl] 0,0-dimethyl phosphorodithioate	OP	insecticide
MCPB	4-(4-chloro-2-methylphenoxy) butyric acid	PH	herbicide
Meta-Systox R	(see oxydemetonmethyl)		
metham sodium	sodium N-methyldithiocarbamate dihydrate	C	fungicide herbicide nematocide
methomyl	S-methyl N-[(methylcarbamoyl)oxy] thioacetimidate	C	insecticide
methoxychlor	1,1,1-trichloro-2,2-bis(p-methoxyphenyl) ethane	OC	insecticide
methyl parathion	0,0-dimethyl O-p-nitrophenyl phosphorothioate	OP	insecticide
mirex	dodecachlorooctahydro-1,3,4-metheno-2H-cyclobuta [cd] pentalene	OC	insecticide

Table 1. Common and chemical names of pesticidal compounds tested in 5-day diets of second-week birds--continued

Common or registered name	Chemical name	Class ^a	Use
Mocap	(see prophos)		
monuron	3-(p-chlorophenyl)-1,1-dimethylurea	U	herbicide
Morsodren	methylmercuric dicyandiamide	OM	fungicide
nabam	disodium ethylene bisdi-thiocarbamate	C	fungicide
naled	1,2-dibromo-2,2-dichloroethyl dimethyl phosphate	OP	insecticide
Ortho 11775	3-(2-butyl)phenyl-N-methyl-N-(phenylsulfenyl) carbamate	C	insecticide
oxydemetonmethyl	S-[2-(ethylsulfinyl)ethyl]0,0-dimethyl phosphorothioate	OP	insecticide
Panogen	(see Morsodren)		
Paraquat CL	(see paraquat dichloride)		
paraquat dichloride	1,1'-dimethyl-4,4'dipyridinium dichloride	O	herbicide
parathion	0,0-diethyl 0-p-nitrophenyl phosphorothioate	OP	acaricide insecticide
Paris green	copper acetoarsenite	OM	insecticide
pentachlorophenol	pentachlorophenol	OC	defoliant herbicide insecticide fungicide wood preserv- ative
Perthane	1,1-dichloro-2,2-bis (p-ethylphenyl) ethane	OC	insecticide

Table 1. Common and chemical names of pesticidal compounds tested in 5-day diets of second-week birds--continued

Common or registered name	Chemical name	Class ^a	Use
phorate	0,0-diethyl S-(ethylthio) methyl phosphorodithioate	OP	insecticide systemic
phosphamidon	2-chloro-2-diethylcarbamoyl-1-methylvinyl dimethyl phosphate	OP	acaricide systemic insecticide
Phygon	(see dichlone)		
picloram	4-amino-3,5,6-trichloropicolinic acid	O	herbicide
prophos	0-ethyl S,S-dipropyl phosphorodithioate	OP	insecticide nematocide
Rhothane	(see DDD)		
Sevin	(see carbaryl)		
silvex	2-(2,4,5-trichlorophenoxy) propionic acid	PH	herbicide
silvex, butoxy-ethanol ester	2-(2,4,5-trichlorophenoxy) propionic acid, butoxyethanol ester	PH	herbicide
simazine	2-chloro-4,6-bis(ethylamino)- <u>s</u> -triazine	O	herbicide
Starlicide	(see DRC-1339)		
Strobane	(see terpene polychlorinates)		
Sumithion	(see fenitrothion)		
Systox	(see demeton)		
2,4,5-T,butoxy-ethanol ester	2,4,5-trichlorophenoxyacetic acid, butoxyethanol ester	PH	herbicide
TDE	(see DDD)		
Tedion	(see tetradifon)		
terpene polychlorinates	chlorinated terpenes	OC	insecticide

Table 1. Common and chemical names of pesticidal compounds tested in 5-day diets of second-week birds--continued

Common or registered name	Chemical name	Class ^a	Use
tetradifon	p-chlorophenyl 2,4,5-trichlorophenyl sulfone	O	insecticide acaricide
TFM	3-trifluoromethyl-4-nitrophenol	O	lampricide
Thimet	(see phorate)		
thionazin	0,0-diethyl 0-2-pyrazinyl phosphorothioate	OP	insecticide nematocide
Thiodan	(see endosulfan)		
thiram	bis (dimethylthiocarbamyl) disulfide	O	fungicide animal repellent
Tordon	(see picloram)		
toxaphene	chlorinated camphene containing 67-69% chlorine	OC	insecticide
2,4,5-TP acid	(see silvex)		
2,4,5-TP,butoxy-ethanol ester	(see silvex, butoxyethanol ester)		
trichlorfon	0,0-dimethyl (1-hydroxy-2,2,2-trichloroethyl) phosphonate	OP	systemic insecticide
Vapam	(see metham sodium)		
Zectran	4-dimethylamino-3,5-xylyl N-methylcarbamate	C	insecticide
Zinophos	(see thionazin)		

^aKey to class designations:

C	carbamate
O	other
OC	organochlorine
OM	organometallic
OP	organophosphate
PH	phenoxy
U	synthetic urea

Table 2 presents toxicity data for each species within an alphabetical listing of chemicals by common name or, as necessary, registered name. Parameters include: the LC_{50} and its 95 percent confidence limits; the slope, with its standard deviation, of the weighted linear regression of probits on log-concentration; and the relative toxicity of the dieldrin standard to the chemical - the "RTD" - and the 95 percent confidence limits of this ratio. The table also shows the number of concentrations and birds per concentration used in each determination, and a "toxic rank number" which, for a given species, is the numerical position of a chemical in a list ranked in descending order of toxicity. No attempt was made to assign toxic rank numbers where LC_{50} 's exceeded 5,000 ppm; however, the percentage of mortality at 5000 ppm is given if deaths occurred.

Table 2. Median lethal concentrations (LC₅₀'s) of pesticidal chemicals in 5-day diets of second-week birds, with relative toxicities of dieldrin (RTD) when tested concurrently as a standard

CHEMICALS		No. concen- trations	(Birds per conc.)	LC ₅₀ : ^a ppm chem. in feed	(95% conf. limits)	Slope: probit on log (St. conc. dev.)	Relative ^b toxicity of dieldrin (RTD)	(95% conf. limits)	Toxic ^c rank no.
Species									
Abate									
Bobwhite	4	(6)	92 (70- 117)	9.842 (2.816)	2.67	(1.94 -3.67)	12	
Jap. quail	6	(16)	260 (206- 334)	5.247 (1.047)	5.27	(3.87 -7.34)	19	
Pheasant	4	(8)	162 (120- 207)	8.135 (3.332)	3.05	(2.22 -4.05)	9	
Mallard	4	(8)	894 (575-1910)	2.739 (1.586)	2.79	(1.42 -4.76)	12	
aldrin									
Bobwhite	6	(10)	37 (33- 41)	9.867 (2.082)	0.943	(0.821-1.09)	5	
Jap. quail	5	(18)	34 (28- 41)	5.133 (1.243)	0.812	(0.663-0.987)	4	
Pheasant	6	(10)	57 (50- 64)	10.433 (1.835)	1.05	(0.875-1.25)	3	
Mallard	6	(10)	155 (129- 186)	4.417 (1.507)	0.763	(0.596-0.978)	4	
amitrole									
Bobwhite	-								
Jap. quail	3	(14)	>5000 ^d						
Pheasant	3	(9)	>5000						
Mallard	3	(10)	>5000						
Aramite									
Bobwhite	3	(10)	>5000	20% mort. @ 5000					
Jap. quail	3	(10)	>5000						
Pheasant	6	(10)	>5000						
Mallard	-								

See footnotes at end of table.

Table 2. Median lethal concentrations (LC₅₀'s) of pesticidal chemicals in 5-day diets of second-week birds, with relative toxicities of dieldrin (RTD) when tested concurrently as a standard--cont.

CHEMICALS									
Species	No. concen- trations	(Birds per conc.)	LC ₅₀ : ^a ppm chem. in feed	(95% conf. limits)	Slope: probit on log conc. dev.)	Relative ^b toxicity of dieldrin (95% conf. (RTD) limits)	Toxic ^c rank no.		
Aroclor 1221									
Bobwhite	6	(10)	>6000	30% mort. @ 6000					
Jap. quail	3	(10)	>6000 ^d						
Pheasant	6	(10)	>4000						
Mallard	-								
Aroclor 1232									
Bobwhite	4	(10)	3002	(2577-3501)	11.631 (2,695)	75.1 (62.0- 92.4)	49		
Jap. quail	3	(10)	>5000						
Pheasant	6	(10)	3146	(2626-3948)	5.786 (1.522)	61.6 ^e -	46		
Mallard	-								
Aroclor 1242									
Bobwhite	5	(10)	2098	(1706-2610)	3.724 (1.739)	70.8 (53.3-101)	48		
Jap. quail	8	(10)	>6000						
Pheasant	5	(10)	2078	(1843-2347)	7.808 (2.616)	40.6 (34.7- 47.7)	42		
Mallard	5	(10)	3182	(2613-3879)	2.577 (1.513)	19.7 (15.0- 26.3)	37		
Aroclor 1248									
Bobwhite	6	(10)	1175	(966-1440)	2.950 (1.355)	39.7 (30.0- 55.8)	40		
Jap. quail	7	(10)	4844	(4355-5410)	7.845 (1,996)	77.4 (66.2- 90.7)	49		
Pheasant	6	(10)	1312	(1166-1477)	7.534 (2.366)	25.7 (21.9- 30.0)	36		
Mallard	5	(10)	2798	(2264-3422)	4.725 (1.516)	17.3 (13.1- 23.0)	33		

Table 2. Median lethal concentrations (LC₅₀'s) of pesticidal chemicals in 5-day diets of second-week birds, with relative toxicities of dieldrin (RTD) when tested concurrently as a standard--cont.

CHEMICALS								
Species	No. concen- trations	(Birds per conc.)	LC50: ^a ppm chem. in feed	(95% conf. limits)	Slope: probit on log (St. dev.)	Relative ^b toxicity of dieldrin (95% conf. limits) (RTD)	Toxic ^c rank no.	
<u>Aroclor 1254</u>								
Bobwhite	5	(10)	604	(410- 840)	6.379	(1.848)	20.4 (15.0 -27.7)	33
Jap. quail	8	(10)	2898	(2598-3241)	5.772	(1.364)	46.3 (39.4 -54.5)	45
Pheasant	5	(10)	1091	(968-1228)	12.174	(2.431)	21.3 (18.2 -25.0)	32
Mallard	6	(10)	2699	(2159-3309)	6.674	(1.263)	16.7 (12.7 -22.0)	31
<u>Aroclor 1260</u>								
Bobwhite	5	(10)	747	(577- 937)	6.211	(1.631)	25.2 (18.9 -34.4)	37
Jap. quail	7	(10)	2186	(1917-2478)	7.444	(1.439)	34.9 (29.3 -41.3)	41
Pheasant	6	(10)	1260	(1106-1433)	5.421	(2.715)	24.6 (20.8 -29.1)	35
Mallard	5	(10)	1975	(1363-2749)	4.054	(1.759)	12.2 (8.93-16.3)	27
<u>Aroclor 1262</u>								
Bobwhite	5	(10)	871	(702-1069)	4.037	(1.584)	29.4 (22.1 -40.8)	39
Jap. quail	7	(10)	2291	(2038-2575)	7.552	(1.501)	36.6 (31.0 -43.2)	43
Pheasant	5	(10)	1234	(1086-1402)	13.518	(2.547)	24.1 (20.5 -28.5)	34
Mallard	6	(10)	3008	(2461-3634)	2.351	(1.226)	18.6 (14.2 -24.5)	36
<u>atrazine</u>								
Bobwhite	3	(10)	>5000 ^d					
Jap. quail	3	(14)	>5000 7% mort. @ 5000					
Pheasant	3	(8)	>5000					
Mallard	3	(10)	>5000 30% mort. @ 5000					

Table 2. Median lethal concentrations (LC₅₀'s) of pesticidal chemicals in 5-day diets of second-week birds, with relative toxicities of dieldrin (RTD) when tested concurrently as a standard--cont.

CHEMICALS		No. (Birds concen- trations conc.)	LC50: ^a ppm chem. in feed	(95% conf. limits)	Slope: probit on log (St. conc. dev.)	Relative ^b toxicity of dieldrin (95% conf. (RTD) limits)	Toxic ^c rank no.
Species							
<u>Baygon</u>							
Bobwhite	4	(10)	206	(168- 251)	4.215 (1.988)	7.68 (5.79-10.6)	19
Jap. quail	—						
Pheasant	—						
Mallard	3	(8)	<1000				
<u>Cadmate</u>							
Bobwhite	5	(10)	1728	(1381-2132)	4.574 (1.162)	41.6 ^e	42
Jap. quail	6	(10)	2693	(2269-3202)	3.671 (1.136)	50.5 (39.7 -64.4)	46
Pheasant	5	(10)	1411	(1202-1657)	4.437 (1.523)	26.9 (21.9 -33.0)	38
Mallard	3	(8)	>5000	10% mort. @ 5000			
<u>captan</u>							
Bobwhite	6	(8)	>2400				
Jap. quail	3	(14)	>5000 ^d				
Pheasant	3	(12)	>5000				
Mallard	3	(10)	>5000				
<u>carbaryl</u>							
Bobwhite	2	(7)	>5000				
Jap. quail	3	(14)	>5000	7% mort. @ 5000			
Pheasant	1	(4)	>5000				
Mallard	4	(6)	>5000				

Table 2. Median lethal concentrations (LC50's) of pesticidal chemicals in 5-day diets of second-week birds, with relative toxicities of dieldrin (RTD) when tested concurrently as a standard--cont.

CHEMICALS		No. concen- trations	(Birds per conc.)	LC50: ^a ppm chem. in feed	(95% conf. limits)	Slope: probit on log (St. conc. dev.)	Relative ^b toxicity of dieldrin (95% conf. limits)		Toxic ^c rank no.
Species							(RTD)		
Ceresan M									
Bobwhite		6	(10)	57	(42- 74)	3.409 (1.671)	1.58	(1.12- 2.31)	8
Jap. quail		6	(10)	100	(84- 118)	7.450 (1.119)	1.87	(1.47- 2.39)	12
Pheasant		6	(10)	146	(127- 167)	5.960 (1.192)	2.15	(1.75- 2.60)	7
Mallard		6	(8)	≈ 50	—		≈ 0.28	—	3
chlordane									
Bobwhite		6	(6)	331	(197- 479)	4.866 (1.760)	7.27	(4.71-10.8)	18
Jap. quail		5	(14)	350	(305- 403)	6.651 (1.220)	5.86	(4.86- 7.08)	21
Pheasant		5	(9)	430	(366- 505)	7.120 (1.775)	8.06	(6.51- 9.94)	17
Mallard		5	(10)	858	(629-1241)	3.796 (1.236)	4.23	(2.99- 6.28)	19
Co-Ral									
Bobwhite		6	(10)	120	(104- 139)	7.348 (1.923)	2.89	(2.36- 3.54)	13
Jap. quail		5	(10)	225	(172- 306)	4.642 (1.049)	≈ 4.00	—	16
Pheasant		6	(10)	318	(277- 364)	7.228 (1.452)	6.06	(5.03- 7.30)	14
Mallard		6	(10)	709	(521-1032)	1.981 (0.993)	3.54	(2.34- 5.53)	16
2,4-D,acetamide									
Bobwhite		—							
Jap. quail		3	(16)	>5000 ^d					
Pheasant		—							
Mallard		—							

Table 2. Median lethal concentrations (LC50's) of pesticidal chemicals in 5-day diets of second-week birds, with relative toxicities of dieldrin (RTD) when tested concurrently as a standard--cont.

CHEMICALS		No. concen- trations	(Birds per conc.)	LC50: ^a ppm chem. in feed	(95% conf. limits)	Slope: probit on log (St. conc. dev.)	Relative ^b toxicity of dieldrin (95% conf. limits)		Toxic ^c rank no.
Species							(RTD)		
2,4-D, butoxy- ethanol ester									
Bobwhite		4	(4)	>5000 ^d					
Jap. quail		4	(14)	>5000					
Pheasant		3	(12)	>5000	17% mort. @ 5000				
Mallard		8	(11)	>5000					
2,4-D, dimethyl- amine salt									
Bobwhite		2	(7)	>5000					
Jap. quail		4	(20)	>5000					
Pheasant		3	(8)	>5000					
Mallard		3	(8)	>5000					
dalapon									
Bobwhite		—							
Jap. quail		3	(14)	>5000					
Pheasant		3	(8)	>5000					
Mallard		3	(10)	>5000					
Dasanit									
Bobwhite		5	(10)	35 (28- 43)		5.076 (3.408)	0.892 (0.705-1.15)		4
Jap. quail		—							
Pheasant		6	(10)	148 (119-179)		5.010 (1.369)	≈ 3.50	—	10
Mallard		6	(10)	43 (36- 51)		5.139 (1.192)	0.210 (0.154-0.264)		2

Table 2. Median lethal concentrations (LC₅₀'s) of pesticidal chemicals in 5-day diets of second-week birds, with relative toxicities of dieldrin (RTD) when tested concurrently as a standard--cont.

CHEMICALS		No.	(Birds concen- trations conc.)	LC ₅₀ : ^a ppm chem. in feed	(95% conf. limits)	Slope: probit on log conc. dev.)	Relative ^b toxicity of dieldrin (95% conf. (RTD) limits)	Toxic ^c rank no.
<u>2,4-DB</u>								
Bobwhite	3	(10)		>5000 ^d	40% mort. @ 5000			
Jap. quail	3	(12)		>5000				
Pheasant	3	(10)		>5000				
Mallard	—							
<u>DDD</u>								
Bobwhite	5	(7)		2178	(1835-2584)	9.379 (2.497)	59.2 (47.7 -74.2)	45
Jap. quail	4	(12)		3165	(2534-3978)	4.613 (1.780)	56.2 (43.0 -74.0)	47
Pheasant	4	(7)		579	(499- 668)	11.956 (3.360)	13.5 (11.1 -16.4)	26
Mallard	6	(10)		4814	(3451-7054)	3.455 (1.343)	24.7 (17.9 -36.1)	40
<u>DDE</u>								
Bobwhite	5	(7)		825	(697- 976)	8.132 (2.436)	22.5 (18.1 -28.0)	35
Jap. quail	6	(12)		1355	(1111-1648)	6.469 (1.205)	24.1 (18.6 -31.0)	36
Pheasant	5	(7)		841	(731- 967)	12.198 (2.969)	19.6 (16.3 -23.7)	29
Mallard	6	(10)		3572	(2811-4669)	3.709 (1.069)	18.4 (13.6 -25.7)	35
<u>DDT</u>								
Bobwhite	5	(7)		611	(514- 724)	7.357 (2.489)	16.6 (13.4 -20.8)	28
Jap. quail	6	(12)		568	(470- 687)	4.770 (1.367)	10.1 (7.86-13.0)	28
Pheasant	4	(7)		311	(256- 374)	10.982 (4.644)	7.27 (5.88- 8.9)	16
Mallard	6	(10)		1869	(1500-2372)	3.896 (0.996)	9.60 (7.14-13.3)	23

Table 2. Median lethal concentrations (LC50's) of pesticidal chemicals in 5-day diets of second-week birds, with relative toxicities of dieldrin (RTD) when tested concurrently as a standard--cont.

CHEMICALS:		No. concen- trations	(Birds per conc.)	LC50: ^a		Slope: probit on log (St. conc. dev.)	Relative ^b toxicity of dieldrin (95% conf. limits)	Toxic ^c rank no.
Species				ppm chem. in feed	(95% conf. limits)			
DDVP								
Bobwhite	-							
Jap. quail	-							
Pheasant	-							
Mallard	3	(10)		>5000	30% mort. @ 5000			
demeton								
Bobwhite	6	(8)		596	(472-768)	4.510 (1.289)	13.5 ^e	24
Jap. quail	6	(10)		275	(228-327)	5.168 (1.314)	5.15	(4.04 - 6.52)
Pheasant	6	(8)		665	(572-773)	7.238 (1.915)	10.2	(8.29 - 12.5)
Mallard	6	(11)		598	(488-733)	2.689 (0.873)	4.28	(3.23 - 5.78)
diazinon								
Bobwhite	4	(8)		245	(178-234)	10.771 (3.271)	6.41	(4.29 - 9.44)
Jap. quail	5	(18)		47	(40- 54)	6.962 (1.017)	1.11	(0.911- 1.35)
Pheasant	4	(8)		244	(177-322)	6.796 (1.794)	6.01	(4.19 - 9.18)
Mallard	5	(10)		191	(138-253)	3.687 (1.186)	0.943	(0.634- 1.33)
dichlobenil								
Bobwhite	-							
Jap. quail	3	(16)		>5000	19% mort. @ 5000			
Pheasant	3	(8)		≈ 1500	-		≈ 27	39
Mallard	-							

Table 2. Meidan lethal concentrations (LC50's) of pesticidal chemicals in 5-day diets of second-week birds, with relative toxicities of dieldrin (RTD) when tested concurrently as a standard--cont.

CHEMICALS		No. concen- trations	(Birds per conc.)	LC50: ^a ppm chem. in feed	(95% conf. limits)	Slope: probit on log conc. dev.)	Relative ^b toxicity of dieldrin (95% conf. limits)		Toxic ^c rank no.
Species									
<u>dichlone</u>									
Bobwhite	2	(10)		>5000 ^d					
Jap. quail	3	(15)		>5000					
Pheasant	3	(8)		>5000					
Mallard	3	(9)		>5000					
<u>dichloro- benzophenone</u>									
Bobwhite	—								
Jap. quail	—								
Pheasant	—								
Mallard	3	(5)		>5000					
<u>dicofof</u>									
Bobwhite	6	(8)		3010	(2635-3424)	4.306	(2.871)	67.9	(56.8-81.0) 47
Jap. quail	6	(14)		1418	(1232-1628)	4.133	(1.002)	26.5	(21.7-32.3) 40
Pheasant	6	(12)		2126	(1892-2387)	7.378	(1.861)	37.1	(31.6-43.6) 41
Mallard	5	(9)		1651	(1356-2029)	5.638	(1.354)	13.7	(10.5-18.8) 29
<u>dieldrin</u>									
Bobwhite	102	(10)		39	(37- 41)	9.631	(0.514)	1	— 6
Jap. quail	78	(10)		57	(53- 61)	6.411	(0.314)	1	— 7
Pheasant	96	(10)		56	(51- 60)	9.930	(0.510)	1	— 2
Mallard	90	(10)		185	(152- 217)	4.722	(0.298)	1	— 6

Table 2. Median lethal concentrations (LC50's) of pesticidal chemicals in 5-day diets of second-week birds, with relative toxicities of dieldrin (RTD) when tested concurrently as a standard--cont.

CHEMICALS		No. concen- trations	(Birds per conc.)	LC50: ^a ppm chem. in feed	Slope: probit on log (St. conc. dev.)	Relative ^b toxicity of dieldrin (95% conf. limits)	Toxic ^c rank no.
Species	(95% conf. limits)						
dimethoate							
Bobwhite	—						
Jap. quail	6 (16)	346 (303- 394)	6.782 (1.273)	5.83 (4.88- 6.95)	20		
Pheasant	6 (8)	332 (293- 376)	10.075 (3.872)	7.05 (5.96- 8.32)	15		
Mallard	6 (9)	1011 (707-1372)	2.017 (0.931)	10.0 (6.54- 17.2)	24		
dioxathion							
Bobwhite	—						
Jap. quail	6 (14)	6640 (5105-9000)	7.195 (1.885)	124 (102 -158)	51		
Pheasant	5 (9)	4067 (3593-4610)	7.769 (2.502)	82.7 (69.1 - 99.3)	50		
Mallard	4 (8)	≈ 3600	—	≈ 18	34		
diquat							
Bobwhite	4 (8)	2932 (1811-5256)	7.588 (2.727)	83.9 (53.6 -145)	50		
Jap. quail	5 (16)	1346 (1178-1540)	4.755 (1.414)	22.7 (19.0 - 27.1)	34		
Pheasant	6 (9)	3742 (3329-4220)	7.507 (2.011)	76.1 (63.9 - 91.1)	49		
Mallard	3 (10)	>5000 33% mort. @ 5000					
disulfoton							
Bobwhite	4 (8)	715 (617- 827)	10.241 (3.310)	16.1 (13.3 - 19.5)	27		
Jap. quail	6 (10)	333 (282- 392)	5.812 (1.244)	6.25 (4.93- 7.91)	23		
Pheasant	5 (9)	634 (547- 737)	7.110 (1.821)	12.9 (10.6 - 15.8)	25		
Mallard	6 (11)	510 (415- 625)	4.713 (0.887)	3.65 (2.75- 4.91)	17		

Table 2. Median lethal concentrations (LC₅₀'s) of pesticidal chemicals in 5-day diets of second-week birds, with relative toxicities of dieldrin (RTD) when tested concurrently as a standard--cont.

CHEMICALS		No. concen- trations	(Birds per conc.)	LC50: ^a ppm chem. in feed	(95% conf. limits)	Slope: probit on log (St. conc. dev.)	Relative ^b toxicity of dieldrin (95% conf. (RTD) limits)	Toxic ^c rank no.
Species								
diuron								
Bobwhite	5	(10)	1730 (1482-2035)	7.218 (1.796)	41.4 (33.8 -51.7)		41	
Jap. quail	3	(14)	>5000 14% mort. @ 5000					
Pheasant	6	(9)	>5000 33% mort. @ 4200; repellent @ 5000					
Mallard	6	(10)	>5000 30% mort. @ 5000					
DRC-1339								
Bobwhite	-							
Jap. quail	6	(13)	23 (20- 26)	7.841 (1.792)	0.394 (0.334- 0.466)		2	
Pheasant	-							
Mallard	-							
Dursban								
Bobwhite	-							
Jap. quail	6	(13)	299 (146-1682)	1.591 (0.766)	5.25 ^e -		18	
Pheasant	-							
Mallard	-							
Dyfonate								
Bobwhite	5	(6)	133 (105- 195)	4.166 (2.764)	3.46 (2.60 - 5.23)		14	
Jap. quail	6	(10)	295 (259- 336)	6.841 (1.476)	6.00 (4.98 - 7.25)		22	
Pheasant	-							
Mallard	5	(10)	1225 (889-1773)	3.399 (1.082)	6.11 (3.96 - 9.59)		22	

Table 2. Median lethal concentrations (LC₅₀'s) of pesticidal chemicals in 5-day diets of second-week birds, with relative toxicities of dieldrin (RTD) when tested concurrently as a standard--cont.

CHEMICALS												
Species	No. concen- trations	(Birds per conc.)	LC ₅₀ : ^a ppm chem. in feed	(95% conf. limits)	Slope: probit on log (St. conc. dev.)	Relative ^b toxicity of dieldrin (RTD)	(95% conf. limits)	Toxic ^c rank no.				
endosulfan												
Bobwhite	5	(8)	805	(690- 939)	4.796	(3.997)	19.9 (16.5 -24.5)	32				
Jap. quail	6	(13)	≈ 1250	—			≈ 22 —	33				
Pheasant	6	(8)	1275	(1098-1482)	5.326	(1,904)	19.6 (15.9 -24.0)	30				
Mallard	4	(10)	1053	(781-1540)	5.316	(1.507)	4.21 (3.01 - 6.26)	18				
endrin												
Bobwhite	6	(10)	14	(11- 24)	2.993	(1.243)	0.370 ^e —	1				
Jap. quail	6	(13)	18	(15- 20)	9.020	(1.844)	0.304 (0.257- 0.359)	1				
Pheasant	4	(8)	14	(11- 17)	3.485	(1.536)	0.335 (0.236- 0.527)	1				
Mallard	6	(10)	22	(17- 31)	3.425	(0.991)	0.108 ^e —	1				
EPN												
Bobwhite	5	(10)	349	(289- 411)	7.547	(2.080)	8.88 (7.10 -11.0)	21				
Jap. quail	5	(10)	443	(349- 550)	3.246	(1.405)	≈ 7.90 —	25				
Pheasant	6	(10)	1075	(943-1230)	6.776	(1.510)	20.5 (17.1 -24.7)	31				
Mallard	3	(5)	< 400									
fenac												
Bobwhite	2	(9)	>5000 ^d									
Jap. quail	3	(16)	>5000									
Pheasant	3	(8)	>5000									
Mallard	3	(9)	>5000									

Table 2. Median lethal concentrations (LC50's) of pesticidal chemicals in 5-day diets of second-week birds, with relative toxicities of dieldrin (RTD) when tested concurrently as a standard--cont.

CHEMICALS		No.	(Birds	LC50: ^a	Slope:	Relative ^b	Toxic ^c
Species	concen- trations	per conc.)	ppm chem.	(95% conf. limits)	probit on log (St. conc. dev.)	dieldrin (95% conf. limits)	rank no.
fenitrothion							
Bobwhite	5	(10)	157	(135- 183)	6.986 (1.936)	3.79 (3.09 - 4.67)	15
Jap. quail	—						
Pheasant	6	(10)	453	(388- 525)	8.131 (2.051)	8.36 (6.89 -10.1)	18
Mallard	5	(10)	2482	(1693-3985)	2.083 (1.166)	12.4 (8.05 -20.2)	28
fenthion							
Bobwhite	5	(8)	30	(21- 41)	6.640 (3.675)	0.784 (0.508- 1.16)	3
Jap. quail	6	(16)	86	(68- 109)	6.361 (0.946)	1.73 (1.28 - 2.37)	11
Pheasant	4	(8)	202	(154- 254)	7.371 (3.071)	3.80 (2.79 - 5.06)	11
Mallard	6	(9)	231	(108- 395)	2.080 (1.115)	2.29 (1.35 - 3.66)	9
fenuron							
Bobwhite	3	(9)	>5000 ^d				
Jap. quail	3	(14)	>5000				
Pheasant	3	(5)	>5000				
Mallard	3	(10)	>5000				
Guthion							
Bobwhite	6	(6)	488	(394- 601)	6.441 (2.395)	14.2 (10.6 -19.4)	25
Jap. quail	6	(16)	639	(512- 796)	4.189 (0.932)	12.9 (9.53 -17.7)	30
Pheasant	4	(8)	1821	(1355-2468)	4.466 (1.504)	44.8 (30.5 -74.0)	43
Mallard	6	(9)	1940	(978-4506)	1.791 (0.587)	11.4 ^e —	25

Table 2. Median lethal concentrations (LC50's) of pesticidal chemicals in 5-day diets of second-week birds, with relative toxicities of dieldrin (RTD) when tested concurrently as a standard--cont.

CHEMICALS		No. concen- trations	(Birds per conc.)	LC50: ^a ppm chem. in feed	(95% conf. limits)	Slope: probit on log conc.	(St. dev.)	Relative ^b toxicity of dieldrin (RTD)	(95% conf. limits)	Toxic ^c rank no.
Species										
<u>heptachlor</u>										
Bobwhite	5	(7)	92	(76- 113)	7.350	(2.233)	2.51	(1.99- 3.22)	10	
Jap. quail	6	(16)	93	(74- 116)	3.722	(0.939)	1.88	(1.39- 2.58)	13	
Pheasant	4	(10)	224	(191- 265)	7.277	(2.876)	4.13	(3.42- 5.04)	12	
Mallard	6	(9)	480	(389- 570)	5.264	(1.646)	2.82	(2.12- 3.71)	13	
<u>Imidan</u>										
Bobwhite	6	(8)	501	(339- 781)	2.422	(0.844)	14.3 ^e	—	26	
Jap. quail	6	(10)	1217	(1065-1392)	4.481	(1.439)	24.7	(20.5 - 30.0)	37	
Pheasant	—									
Mallard	3	(8)	>5000	40% mort. @ 5000						
<u>lindane</u>										
Bobwhite	6	(10)	882	(755-1041)	2.456	(1.673)	21.1	(17.2 - 26.4)	34	
Jap. quail	6	(15)	425	(347- 520)	3.487	(0.692)	7.12 ^e	—	24	
Pheasant	5	(8)	561	(445- 690)	8.251	(2.752)	10.6	(7.88- 14.1)	23	
Mallard	6	(12)	>5000	40% mort. @ 5000						
<u>malathion</u>										
Bobwhite	6	(6)	3497	(2959-4117)	5.931	(2.533)	102	(78.2 -135)	51	
Jap. quail	6	(12)	2128	(1780-2546)	3.622	(1.278)	37.8	(29.7 - 48.2)	44	
Pheasant	5	(7)	4320	(3847-4886)	6.736	(3.423)	101	(85.5 -120)	53	
Mallard	3	(10)	>5000 ^d							

Table 2. Median lethal concentrations (LC50's) of pesticidal chemicals in 5-day diets of second-week birds, with relative toxicities of dieldrin (RTD) when tested concurrently as a standard--cont.

CHEMICALS							
Species	No. concen- trations	(Birds per conc.)	LC ₅₀ : ^a ppm chem. in feed	(95% conf. limits)	Slope: probit on log (St. conc. dev.)	Relative ^b toxicity of dieldrin (95% conf. (RTD) limits)	Toxic ^c rank no.
MCPB							
Bobwhite	3	(10)	>5000	10% mort. @ 5000			
Jap. quail	3	(12)	>5000 ^d				
Pheasant	3	(10)	>5000				
Mallard	—						
metham sodium							
Bobwhite	2	(10)	>5000				
Jap. quail	3	(14)	>5000	12% mort. @ 5000			
Pheasant	3	(8)	>5000				
Mallard	2	(8)	>5000				
methomyl							
Bobwhite	4	(10)	≈ 1100			≈ 28	38
Jap. quail	6	(10)	3124	(2513-3940)	2.682 (2.147)	59.5 (46.7 -77.0)	48
Pheasant	—						
Mallard	6	(10)	2883	(2000-4572)	1.283 (1.086)	16.7 (9.69-31.7)	32
methoxychlor							
Bobwhite	2	(7)	>5000				
Jap. quail	3	(12)	>5000				
Pheasant	3	(5)	>5000				
Mallard	3	(10)	>5000				

Table 2. Median lethal concentrations (LC₅₀'s) of pesticidal chemicals in 5-day diets of second-week birds, with relative toxicities of dieldrin (RTD) when tested concurrently as a standard--cont.

CHEMICALS		No. concen- trations	(Birds per conc.)	LC50: ^a		Slope: probit on log (St. conc. dev.)	Relative ^b toxicity of dieldrin (95% conf. limits)		Toxic ^c rank no.
Species				ppm chem. in feed	(95% conf. limits)		(RTD)		
<u>methyl parathion</u>									
Bobwhite	6	(6)	90 (73- 111)	5.240 (2.164)	2.63 (1.96 - 3.57)		11		
Jap. quail	6	(12)	46 (38- 55)	5.395 (1.100)	0.816 (0.643- 1.03)		5		
Pheasant	6	(7)	116 (101- 134)	11.975 (2.935)	2.71 (2.24 - 3.28)		8		
Mallard	5	(10)	682 (541- 892)	3.216 (1.227)	4.98 (3.61 - 7.83)		21		
<u>mirex</u>									
Bobwhite	6	(10)	2511 (2160-2908)	6.032 (1.731)	60.6 (49.3 -74.2)		46		
Jap. quail	3	(10)	>5000 20% mort. @ 5000						
Pheasant	6	(10)	1540 ^d (1320-1789)	5.801 (1.508)	29.3 (24.1 -35.7)		40		
Mallard	3	(8)	>5000 ^d						
<u>monuron</u>									
Bobwhite	3	(6)	>5000						
Jap. quail	3	(14)	>5000 21% mort. @ 5000						
Pheasant	5	(9)	4682 (3902-5746)	8.213 (2.470)	87.7 (72.1 -109)		51		
Mallard	3	(10)	>5000 10% mort. @ 5000						
<u>Morsodren</u>									
Bobwhite	-								
Jap. quail	6	(10)	47 (40- 56)	7.745 (1.466)	0.964 (0.784- 1.19)		6		
Pheasant	-								
Mallard	-								

Table 2. Median lethal concentrations (LC₅₀'s) of pesticidal chemicals in 5-day diets of second-week birds, with relative toxicities of dieldrin (RTD) when tested concurrently as a standard--cont.

CHEMICALS		LC ₅₀ : ^a		Slope:		Relative ^b		Toxic ^c
Species	No. (Birds per concn- trations conc.)	ppm chem. in feed	(95% conf. limits)	probit on log conc. dev.)	dieldrin (RTD) (95% conf. limits)	of dieldrin (RTD) limits)	rank no.	
nabam								
Bobwhite	2 (9)	>5000	11% mort. @ 5000					
Jap. quail	3 (15)	>5000 ^d						
Pheasant	3 (8)	>5000						
Mallard	3 (10)	>5000						
naled								
Bobwhite	6 (8)	2117	(1502- 2890)	5.169	(3.257)	55.5	(36.4-83.8)	44
Jap. quail	5 (20)	1327	(1178- 1490)	6.542	(1.059)	23.3	(19.9-27.3)	35
Pheasant	5 (10)	2538	(2221- 2896)	4.905	(1.974)	46.8	(39.0-56.2)	44
Mallard	5 (10)	2724	(1068-15089)	0.912	(0.792)	16.0 ^e	—	30
Ortho 11775								
Bobwhite	5 (8)	1474	(1075- 2108)	8.368	(1.814)	42.2	(27.2-68.0)	43
Jap. quail	6 (10)	1345	(1139- 1588)	7.810	(1.366)	25.6	(20.5-32.1)	39
Pheasant	6 (10)	2874	(2567- 3209)	9.888	(2.273)	51.7	(44.4-60.2)	45
Mallard	3 (8)	≈ 2300	—			≈ 11	—	26
oxydemetonmethyl								
Bobwhite	5 (8)	434	(304- 600)	5.209	(1.714)	12.4	(7.67-19.4)	23
Jap. quail	6 (10)	1309	(1097- 1552)	4.163	(1.352)	25.0	(19.8-31.3)	38
Pheasant	6 (10)	1497	(1326- 1690)	9.292	(2.412)	25.7	(22.0-29.9)	37
Mallard	3 (8)	>5000	40% mort. @ 5000					

Table 2. Median lethal concentrations (LC₅₀'s) of pesticidal chemicals in 5-day diets of second-week birds, with relative toxicities of dieldrin (RTD) when tested concurrently as a standard--cont.

CHEMICALS								
Species	No. concen- trations	(Birds per conc.)	LC ₅₀ : ^a ppm chem. in feed	(95% conf. limits)	Slope: probit on log (St. conc. dev.)	Relative ^b toxicity of dieldrin (95% conf. (RTD) limits)	Toxic ^c rank no.	
<u>paraquat dichloride</u>								
Bobwhite	6	(10)	981	(784-1213)	5.022 (1.283)	25.0 ^e —	36	
Jap. quail	6	(10)	970	(823-1140)	6.059 (1.307)	18.5 (14.8 -23.1)	31	
Pheasant	6	(10)	1468	(1287-1673)	5.846 (1.973)	22.3 (18.7 -26.5)	33	
Mallard	6	(10)	4048	(3432-4886)	6.765 (1.281)	23.5 (18.5 -30.7)	38	
<u>parathion</u>								
Bobwhite	5	(6)	194	(150- 245)	4.690 (2.636)	5.65 (4.15 - 7.68)	16	
Jap. quail	6	(12)	44	(36- 53)	5.370 (1.212)	0.775 (0.607- 0.985)	3	
Pheasant	5	(7)	365	(316- 420)	11.039 (3.047)	8.52 (7.04 -10.3)	19	
Mallard	5	(10)	275	(183- 373)	4.383 (1.375)	2.01 (1.40 - 2.86)	8	
<u>Paris green</u>								
Bobwhite	6	(8)	480	(206-2042)	3.474 (1.920)	10.9 ^e —	22	
Jap. quail	5	(20)	1204	(1069-1351)	4.925 (1.105)	21.1 (18.0 -24.7)	32	
Pheasant	6	(8)	1043	(896-1217)	6.644 (2.043)	16.0 (13.0 -19.7)	27	
Mallard	6	(10)	>5000 20% mort. @ 5000					
<u>pentachlorophenol</u>								
Bobwhite	—							
Jap. quail	5	(16)	5204	(4536-6034)	6.877 (1.790)	87.6 (73.9 -105)	50	
Pheasant	6	(12)	4331	(3926-4787)	8.990 (1.945)	75.5 (65.2 - 88.0)	48	
Mallard	2	(8)	≈ 4500	—		≈ 24 —	39	

Table 2. Median lethal concentrations (LC₅₀'s) of pesticidal chemicals in 5-day diets of second-week birds, with relative toxicities of dieldrin (RTD) when tested concurrently as a standard--cont.

CHEMICALS									
Species	No. concen- trations per (Birds conc.)	LC ₅₀ : ^a ppm chem. in feed	(95% conf. limits)	Slope: probit on log (St. conc. dev.)	Relative toxicity of dieldrin (RTD)	(95% conf. limits)	Toxic rank no.		
Perthane									
Bobwhite	3 (10)	>5000	10% mort. @ 5000						
Jap. quail	4 (20)	>5000 ^d							
Pheasant	3 (8)	>5000							
Mallard	3 (10)	>5000							
phorate									
Bobwhite	4 (8)	373	(326- 431)	16.173 (3.640)	8.43	(7.04 -10.2)	20		
Jap. quail	3 (15)	≈ 200	—		≈ 3.6	—	15		
Pheasant	6 (9)	441	(381- 510)	7.648 (1.693)	8.97	(7.36 -10.9)	20		
Mallard	6 (11)	248	(198- 306)	4.853 (0.924)	1.77	(1.33 - 2.38)	7		
phosphamidon									
Bobwhite	6 (8)	24	(10- 37)	3.691 (1.451)	0.632 ^e	—	2		
Jap. quail	6 (14)	89	(77- 102)	5.818 (1.057)	1.16	(0.937- 1.41)	9		
Pheasant	6 (10)	77	(68- 87)	6.564 (1.809)	1.42	(1.19 - 1.70)	5		
Mallard	6 (10)	712	(558- 887)	3.860 (1.154)	3.51	(2.47 - 4.82)	15		
picloram									
Bobwhite	3 (10)	>5000							
Jap. quail	3 (14)	>5000							
Pheasant	3 (8)	>5000							
Mallard	3 (10)	>5000							

Table 2. Median lethal concentrations (LC₅₀'s) of pesticidal chemicals in 5-day diets of second-week birds, with relative toxicities of dieldrin (RTD) when tested concurrently as a standard--cont.

CHEMICALS		^a		^b		^c	
Species	No. (Birds concentrations conc.)	LC50: ppm chem. in feed	(95% conf. limits)	Slope: probit on log (St. conc. dev.)	Relative toxicity of dieldrin (95% conf. limits) (RTD)	rank	Toxic no.
prophos							
Bobwhite	6 (10)	33 (27- 40)	4.956 (1.437)	1.24 (0.947-	1.66)	7
Jap. quail	6 (10)	100 (85- 117)	5.285 (1.498)	2.03 (1.65 -	2.50)	14
Pheasant	6 (10)	118 (103- 134)	9.500 (1.956)	1.79 (1.50 -	2.12)	6
Mallard	3 (8)	<1000					
silvex							
Bobwhite	—	>5000 ^d					
Jap. quail	3 (14)	≈ 4500	—		≈ 96	—	52
Pheasant	3 (8)						
Mallard	—						
silvex, butoxy- ethanol ester							
Bobwhite	3 (10)	3031	(2441-3774)	10.808 (3.945)	113	(84.2 -160)	52
Jap. quail	3 (16)	>5000	12% mort. @ 5000				
Pheasant	3 (8)	<3000					
Mallard	2 (8)	>5000					
simazine							
Bobwhite	3 (10)	>5000					
Jap. quail	3 (14)	>3720					
Pheasant	3 (8)	>5000					
Mallard	3 (9)	>5000					

Table 2. Median lethal concentrations (LC₅₀'s) of pesticidal chemicals in 5-day diets of second-week birds, with relative toxicities of dieldrin (RTD) when tested concurrently as a standard--cont.

CHEMICALS		No. concen- trations	(Birds per conc.)	LC50: ^a ppm chem. in feed	(95% conf. limits)	Slope: probit on log conc. dev.)	Relative toxicity of dieldrin (95% conf. limits)		Toxic rank no. ^c
Species							(RTD)		
2,4,5-T, butoxy- ethanol ester									
Bobwhite	-			>5000 ^d					
Jap. quail	3	(15)		3950 (3106-6118)	4.939 (1.901)		67.8 ^e	-	47
Pheasant	5	(10)		>5000	10% mort. @ 5000				
Mallard	3	(10)							
terepene poly- chlorinates									
Bobwhite	5	(10)		817 (686- 976)	8.311 (1.719)		19.6	(15.7 -24.7)	31
Jap. quail	6	(13)		681 (541- 917)	3.614 (0.938)		11.9 ^e	-	29
Pheasant	6	(8)		800 (698- 920)	10.048 (1.918)		12.3	(10.1 -15.0)	24
Mallard	5	(11)		451 (362- 557)	5.894 (1.169)		3.23	(2.42- 4.34)	14
tetradifon									
Bobwhite	3	(10)		>5000	10% mort. @ 5000				
Jap. quail	3	(14)		>5000					
Pheasant	3	(8)		>5000					
Mallard	3	(9)		>5000					
TFM									
Bobwhite	-								
Jap. quail	-								
Pheasant	-								
Mallard	3	(10)		>5000					

Table 2. Median lethal concentrations (LC50's) of pesticidal chemicals in 5-day diets of second-week birds, with relative toxicities of dieldrin (RTD) when tested concurrently as a standard--cont.

CHEMICALS							
Species	No. concen- trations	(Birds per conc.)	LC50: ^a ppm chem. in feed	(95% conf. limits)	Slope: probit on log (St. conc. dev.)	Relative ^b toxicity of dieldrin (95% conf. (RTD) limits)	Toxic ^c rank no.
thionazin							
Bobwhite	5	(10)	65	(53- 78)	3.520 (1.465)	2.42 (1.83 - 3.35)	9
Jap. quail	6	(10)	58	(49- 68)	8.316 (1.431)	1.18 (0.958- 1.45)	10
Pheasant	6	(10)	72	(63- 82)	5.148 (2.593)	1.30 (1.10 - 1.54)	4
Mallard	—						
thiram							
Bobwhite	—						
Jap. quail	3	(10)	>5000 ^d				
Pheasant	4	(10)	>5000				
Mallard	2	(9)	>5000	20% mort. @ 5000			
toxaphene							
Bobwhite	6	(6)	828	(619-1102)	2.509 (1.406)	18.2 ^e (12.6 -29.1)	29
Jap. quail	6	(14)	686	(523-1002)	2.796 (0.782)	8.72 ^e —	26
Pheasant	5	(9)	542	(462- 638)	5.917 (1.734)	10.2 ^e (8.22 -12.5)	21
Mallard	4	(8)	538	(474- 614)	14.113 (3.128)	2.6 ^e —	11

Table 2. Median lethal concentrations (LC₅₀'s) of pesticidal chemicals in 5-day diets of second-week birds, with relative toxicities of dieldrin (RTD) when tested concurrently as a standard--cont.

CHEMICALS		^a		^b		Toxic ^c rank no.
No. concen- trations	(Birds per conc.)	LC ₅₀ : ppm chem. in feed	(95% conf. limits)	Slope: probit on log (St. conc. dev.)	Relative toxicity of dieldrin (95% conf. limits)	
<u>trichlorfon</u>						
Bobwhite	5	(10)	720 (591- 871)	5.604 (2.677)	18.3 (14.5 -23.3)	30
Jap. quail	6	(10)	1901 (1601-2255)	4.898 (1.108)	35.6 (28.0 -45.4)	42
Pheasant	—					
Mallard	—					
<u>Zectran</u>						
Bobwhite	—				≈ 8.9	27
Jap. quail	3	(14)	≈ 500	—	17.2 (14.0 -21.0)	28
Pheasant	5	(9)	846 (724- 985)	6.558 (1.936)	2.39 (1.79- 3.21)	10
Mallard	6	(11)	334 (268- 412)	3.041 (0.921)		

^aLC₅₀: ppm chemical in ad libitum diet expected to produce 50% mortality in 8 days comprising 5 days of toxic diet followed by 3 days of untreated diet.

^bRelative toxicity of dieldrin (RTD) read: "Dieldrin is x times as toxic as the given chemical as tested." See text for use of RTD's to compare toxicities of any two chemicals.

^cToxic rank number: Numerical position in listing of chemicals ranked in descending order of toxicity (see table 3). The larger the toxic rank number or the RTD, the less toxic the chemical.

^dNo mortality at 5000 ppm unless specified.

^eRelative toxicity of dieldrin applies only at LC₅₀, since probit slope is significantly different (P=0.05) from that of dieldrin.

Table 3a lists chemicals in descending order of toxicity (i.e., by toxic rank number) for each species. The order of toxicity is not based on comparisons of LC₅₀ values *per se* but on RTD values, so as to adjust comparison for possible differences in sensitivity among tests conducted at different times. RTD values and toxic rank numbers are common to Tables 2 and 3a, the latter serving to cross-index the two tables.

Table 3b is an alphabetical listing, by species, of those chemicals having LC₅₀'s exceeding 5,000 ppm or otherwise not known precisely.

Table 3a. Numerical listing of test chemicals ranked in descending order of toxicity, as determined by comparing relative toxicities of a concurrent dieldrin standard

Toxic ^c rank no.	Bobwhite		Japanese quail		Pheasant		Mallard	
	Chemical	RTD ^b	Chemical	RTD	Chemical	RTD	Chemical	RTD
1	endrin	0.370 ^e	endrin	0.304	endrin	0.335	endrin	0.108 ^e
2	phosphamidon	0.632 ^e	DRG-1339	0.394	dieldrin	1.00	Dasanit	0.210
3	fenthion	0.784	parathion	0.775	aldrin	1.05	Ceresan M	≈ 0.28
4	Dasanit	0.892	aldrin	0.812	thionazin	1.30	aldrin	0.763
5	aldrin	0.943	methyl parathion	0.816	phosphamidon	1.42	diazinon	0.943
6	dieldrin	1.00	Morsodren	0.964	prophos	1.79	dieldrin	1.00
7	prophos	1.24	dieldrin	1.00	Ceresan M	2.15	phorate	1.77
8	Ceresan M	1.58	diazinon	1.11	methyl parathion	2.71	parathion	2.01
9	thionazin	2.42	phosphamidon	1.16	Abate	3.05	fenthion	2.29
10	heptachlor	2.51	thionazin	1.18	Dasanit	≈ 3.50	Zectran	2.39
11	methyl parathion	2.63	fenthion	1.73	fenthion	3.80	toxaphene	2.6 ^e
12	Abate	2.67	Ceresan M	1.87	heptachlor	4.13	Abate	2.79
13	Co-Ral	2.89	heptachlor	1.88	diazinon	6.01	heptachlor	2.82
14	Dyfonate	3.46	prophos	2.03	Co-Ral	6.06	terepene polychlorinates	3.23

Table 3a. Numerical listing of test chemicals ranked in descending order of toxicity, as determined by comparing relative toxicities of a concurrent dieldrin standard--continued

Toxic ^c rank no.	Bobwhite		Japanese quail		Pheasant		Mallard	
	Chemical	RTD ^b	Chemical	RTD	Chemical	RTD	Chemical	RTD
15	fenitrothion	3.79	phorate	≈ 3.6	dimethoate	7.05	phosphamidon	3.51
16	parathion	5.65	Co-Ral	≈ 4.00	DDT	7.27	Co-Ral	3.54
17	diazinon	6.41	demeton	5.15	chlordan	8.06	disulfoton	3.65
18	chlordan	7.27	Dursban	5.25 ^e	fenitrothion	8.36	endosulfan	4.21
19	Baygon	7.68	Abate	5.27	parathion	8.52	chlordan	4.23
20	phorate	8.43	dimethoate	5.83	phorate	8.97	demeton	4.28
21	EPN	8.88	chlordan	5.86	toxaphene	10.2	methyl parathion	4.98
22	Paris green	10.9 ^e	Dyfonate	6.00	demeton	10.2	Dyfonate	6.11
23	oxydemeton- methyl	12.4	disulfoton	6.25	lindane	10.6	DDT	9.60
24	demeton	13.5 ^e	lindane	7.12 ^e	terepene polychlorinates	12.3	dimethoate	10.0
25	Guthion	14.2	EPN	≈ 7.90	disulfoton	12.9	Guthion	11.4 ^e
26	Imidan	14.3 ^e	toxaphene	8.72 ^e	DDD	13.5	Ortho 11775	≈ 11
27	disulfoton	16.1	Zectran	≈ 8.9	Paris green	16.0	Aroclor 1260	12.2

Table 3a. Numerical listing of test chemicals ranked in descending order of toxicity, as determined by comparing relative toxicities of a concurrent dieldrin standard--continued

Toxic rank no.	Bobwhite		Japanese quail		Pheasant		Mallard	
	Chemical	RTD ^b	Chemical	RTD	Chemical	RTD	Chemical	RTD
28	DDT	16.6	DDT	10.1	Zectran	17.2	fenitrothion	12.4
29	toxaphene	18.2	terepene polychlorinates	11.9 ^e	DDE	19.6	dicofol	13.7
30	trichlorfon	18.3	Guthion	12.9	endosulfan	19.6	naled	16.0 ^e
31	terepene polychlorinates	19.6	paraquat dichloride	18.5	EPN	20.5	Arocior 1254	16.7
32	endosulfan	19.9	Paris green	21.1	Arocior 1254	21.3	methomyl	16.7
33	Arocior 1254	20.4	endosulfan	≈ 22	paraquat dichloride	22.3	Arocior 1248	17.3
34	lindane	21.1	diquat	22.7	Arocior 1262	24.1	dioxathion	≈ 18
35	DDE	22.5	naled	23.3	Arocior 1260	24.6	DDE	18.4
36	paraquat dichloride	25.0 ^e	DDE	24.1	Arocior 1248	25.7	Arocior 1262	18.6
37	Arocior 1260	25.2	Imidan	24.7	oxydemeton- methyl	25.7	Arocior 1242	19.7
38	methomyl	≈ 28	oxydemeton- methyl	25.0	Cadmate	26.9	paraquat dichloride	23.5
39	Arocior 1262	29.4	Ortho 11775	25.6	dichlobenil	≈ 27	pentachloro- phenol	≈ 24

Table 3a. Numerical listing of test chemicals ranked in descending order of toxicity, as determined by comparing relative toxicities of a concurrent dieldrin standard--continued

Toxic rank no.	Bobwhite		Japanese quail		Pheasant		Mallard	
	Chemical	RTD ^b	Chemical	RTD	Chemical	RTD	Chemical	RTD
40	Aroclor 1248	39.7	dicofo1	26.5	mirex	29.3	DDD	24.7
41	diuron	41.4	Aroclor 1260	34.9	dicofo1	37.1		
42	Cadmate	41.6 ^e	trichlorfon	35.6	Aroclor 1242	40.6		
43	Ortho 11775	42.2	Aroclor 1262	36.6	Guthion	44.8		
44	naled	55.5	malathion	37.8	naled	46.8		
45	DDD	59.2	Aroclor 1254	46.3	Ortho 11775	51.7		
46	mirex	60.6	Cadmate	50.5	Aroclor 1232	61.6 ^e		
47	dicofo1	67.9	DDD	56.2	2,4,5-T, butoxyethanol ester	67.8 ^e		
48	Aroclor 1242	70.8	methomyl	59.5	pentachloro- phenol	75.5		
49	Aroclor 1232	75.1	Aroclor 1248	77.4	diquat	76.1		
50	diquat	83.9	pentachloro- phenol	87.6	dioxathion	82.7		
51	malathion	102	dioxathion	124	monuron	87.7		
52	silvex, butoxyethanol ester	113			silvex	≈ 96		
53					malathion	101		

See footnotes at end of table 2.

Table 3b. Alphabetical listing of chemicals not assigned toxic rank numbers because LC₅₀'s exceeded 5000 ppm or were otherwise not known precisely. Unlisted RTD (LC₅₀'s > 5000 ppm) exceed 128, 87, 90, or 27 for bobwhite, Japanese quail, pheasants, or mallards

Chemical	Bobwhite		Japanese quail		Pheasant		Mallard	
	Chemical	RTD	Chemical	RTD	Chemical	RTD	Chemical	RTD
Aramite			amitrole		amitrole		amitrole	
Aroclor 1221			Aramite		Aramite		atrazine	
atrazine			Aroclor 1221		Aroclor 1221		Baygon	< 5.4
captan	> 62		Aroclor 1232		atrazine		Cadmate	
carbaryl			Aroclor 1242		captan		captan	
2,4-D, butoxy- ethanol ester			atrazine		carbaryl		carbaryl	
2,4-D, dimethyl- amine salt			captan		2,4-D, butoxy- ethanol ester		2,4-D, butoxy- ethanol ester	
2,4-DB			carbaryl		2,4-D, dimethyl- amine salt		2,4-D, dimethyl- amine salt	
dichlone			2,4-D, acetamide		dalapon		dalapon	
fenac			2,4-D, butoxy- ethanol ester		2,4-DB		DDVP	
fenuron			2,4-D, dimethyl- amine salt		dichlone		dichlone	
MCPB			dalapon		diuron		dichloro- benzophenone	

Table 3b. Alphabetical listing of chemicals not assigned toxic rank numbers because LC₅₀'s exceeded 5000 ppm or were otherwise not known precisely. Unlisted RTD (LC₅₀'s >5000 ppm) exceed 128, 87, 90, or 27 for bobwhite, Japanese quail, pheasants, or mallards--continued

Chemical	Bobwhite ^b		Japanese quail		Pheasant		Mallard	
	Chemical	RTD	Chemical	RTD	Chemical	RTD	Chemical	RTD
metham sodium			2,4-DB		fenac		diquat	
metnoxychlor			dichlone		fenuron		diuron	
monuron			dichlobenil		MCPB		EPN	< 2.2
nabam			diuron		metham sodium		fenac	
Perthane			fenac		methoxychlor		fenuron	
picloram			fenuron		nabam		Imidan	
simazine			MCPB		Perthane		lindane	
tetradifon			metham sodium		picloram		malathion	
			methoxychlor		silvex, butoxy- ethanol ester	≈ 36	metham sodium	
			mirex		simazine		methoxychlor	
			monuron		tetradifon		mirex	
			nabam		thiram		monuron	
			Perthane				nabam	
			picloram				oxydemeton- methyl	

Table 3b. Alphabetical listing of chemicals not assigned toxic rank numbers because LC₅₀'s exceeded 5000 ppm or were otherwise not known precisely. Unlisted RTD (LC₅₀'s >5000 ppm) exceed 128, 87, 90, or 27 for bobwhite, Japanese quail, pheasants, or mallards--continued

Bobwhite Chemical	RTD	Japanese quail		Pheasant		Mallard	
		Chemical	RTD	Chemical	RTD	Chemical	RTD
		silvex				Paris green	
		silvex, butoxy- ethanol ester				Perthane	
		simazine	> 65			picloram	
		2,4,5-T, butoxy- ethanol ester				prophos	< 5.4
		tetradifon				silvex, butoxy- ethanol ester	
		thiram				simazine	
						2,4,5-T, butoxy- ethanol ester	
						tetradifon	
						TFM	
						thiram	

See footnotes at end of table 2.

Table 4 provides gross comparisons of species sensitivity based on those chemicals tested on all four species and having LC₅₀'s below 5,000 ppm for at least three species. Findings are presented as percentages of compounds to which each species was the most sensitive, secondmost sensitive, thirdmost sensitive, and least sensitive of the four species. Separate evaluations are presented for organochlorines, organophosphates, polychlorinated biphenyls, and miscellaneous compounds.

Table 4. Percentages of chemicals to which each species was the most sensitive, secondmost, thirdmost, and least sensitive of the four species when chemicals were mixed in 5-day diets (Where ties occurred, percentages were divided equally.)

Sensitivity	Species			
	Bobwhite	Jap. quail	Pheasant	Mallard
<u>(14 organochlorines)^a</u>				
Most	36	25	28	11
2nd most	29	25	28	18
3rd most	14	46	40	0
Least	<u>21</u>	<u>4</u>	<u>4</u>	<u>71</u>
Total	100	100	100	100
<u>(15 organophosphates)^b</u>				
Most	47	53	0	0
2nd most	30	34	13	23
3rd most	13	13	64	10
Least	<u>10</u>	<u>0</u>	<u>23</u>	<u>67</u>
Total	100	100	100	100
<u>(5 Aroclors)^c</u>				
Most	90	0	10	0
2nd most	10	0	90	0
3rd most	0	20	0	80
Least	<u>0</u>	<u>80</u>	<u>0</u>	<u>20</u>
Total	100	100	100	100
<u>(5 other compounds)^d</u>				
Most	40	30	20	10
2nd most	60	10	20	10
3rd most	0	60	40	0
Least	<u>0</u>	<u>0</u>	<u>20</u>	<u>80</u>
Total	100	100	100	100

^a aldrin	^b Abate	^c Aroclor 1242	^d Cadminate
chlordane	Co-Ral	Aroclor 1248	Ceresan M
DDD	demeton	Aroclor 1254	diquat
DDE	diazinon	Aroclor 1260	paraquat
DDT	disulfoton	Aroclor 1262	Paris green
dieldrin	fenthion		
dicofol	Guthion		
endosulfan	malathion		
endrin	m. parathion		
heptachlor	naled		
lindane	oxydemetonmethyl		
mirex	parathion		
terepene	phorate		
polychlorinates	phosphamidon		
toxaphene	prophos		

APPENDIX 1

Calculation of Relative Toxicity Using RTD Values

The RTD values listed in Table 2 can be used to calculate the relative toxicity of two chemicals for a particular species, as follows:

1. Compute the relative toxicity of "Chemical 1" to "Chemical 2" by dividing the RTD of Chemical 2 by the RTD of Chemical 1. For example, if dieldrin is four times as toxic as Chemical 1 ($RTD_1 = 4$) and six times as toxic as Chemical 2 ($RTD_2 = 6$), then Chemical 1 is 1.5 times as toxic as Chemical 2 (i.e., $RTD_2/RTD_1 = 6/4 = 1.5$). An algebraic argument for the procedure is developed in Appendix 2. The calculation of confidence limits for relative toxicities requires more data than could reasonably be included in this report.
2. Test the slopes of the probit regression lines of the two chemicals for parallelism using a 2-tailed t-test. Let b_1 and b_2 be the estimated slopes and s_1 and s_2 their standard deviations. (The s values are actually in standard error form.) Also let n_1 and n_2 equal the number of concentrations used in the respective determinations.

Then

$$t = (b_1 - b_2) / \sqrt{s_1^2 + s_2^2}.$$

Since s_1^2 and s_2^2 have $n_1 - 2$ and $n_2 - 2$ degrees of freedom, t is given $n_1 + n_2 - 4$ degrees of freedom, provided s_1^2 and s_2^2 can be considered estimates of a common σ^2 . We expect homogeneity of variances in most instances; however, procedures for testing the equality of two variances and the significance of the difference of two means (i.e., b_1 and b_2) when variances are unequal are presented in Snedecor and Cochran (1967).

APPENDIX 2

Theoretical Estimation of Relative Toxicity Using RTD Values

We have defined the RTD as the relative toxicity of a dieldrin standard to a given chemical, as derived from their LC_{50} . The use of RTD's to estimate the relative toxicity of two chemicals tested in different experiments is based on the following argument:

Let

C_1 = the true LC_{50} of chemical 1,

C_2 = the true LC_{50} of chemical 2,

S = the true LC_{50} of the standard,

c_1 = the estimated LC_{50} of chemical 1 from experiment I,

c_2 = the estimated LC_{50} of chemical 2 from experiment II,

s_1 = the estimated LC_{50} of the standard from experiment I and,

s_2 = the estimated LC_{50} of the standard from experiment II.

Also, assume f_1 and f_2 are sensitivity factors operative in experiments I and II, such that

$$f_1 s_1 = f_2 s_2 = S,$$

$$f_1 c_1 = C_1, \text{ and}$$

$$f_2 c_2 = C_2$$

Then

$$\text{"RTD}_1\text{"} = f_1 c_1 / f_1 s_1 = C_1 / S, \text{ and}$$

$$\text{"RTD}_2\text{"} = f_2 c_2 / f_2 s_2 = C_2 / S$$


Therefore, the true relative toxicity of chemical 1 to chemical 2 can be estimated as $\text{RTD}_2 / \text{RTD}_1$, since

$$\text{RTD}_2 / \text{RTD}_1 = (C_2 / S) / (C_1 / S) = C_2 / C_1,$$

the true relative toxicity of chemical 1 to chemical 2.

As the Nation's principal conservation agency, the Department of the Interior has basic responsibilities for water, fish, wildlife, mineral, land, park, and recreational resources. Indian and Territorial affairs are other major concerns of this department of natural resources.

The Department works to assure the wisest choice in managing all our resources so that each shall make its full contribution to a better United States now and in the future.



UNITED STATES
DEPARTMENT OF THE INTERIOR
FISH AND WILDLIFE SERVICE
BUREAU OF SPORT FISHERIES AND WILDLIFE
WASHINGTON, D. C. 20240

POSTAGE AND FEES PAID
U.S. DEPARTMENT OF THE INTERIOR

